arXiv:1607.08306v2 [math.DG] 1 Aug 2016

Partial result of Yau's Conjecture of the first eigenvalue in unit sphere $\mathbb{S}^{n+1}(1)$

Zhongyang Sun

Abstract : In this paper, we partially solve Yau' Conjecture of the first eigenvalue of an embedded compact minimal hypersurface of unit sphere $\mathbb{S}^{n+1}(1)$, i.e., Corollary 1.2. In particular, Corollary 1.3 proves that the condition $\int_{\Omega_1}|\nabla u|^2=(n+1)\int_{\Omega_1}u^2$ is naturally true and meaningful in Corollary 1.2.

Keywords : Minimal hypersurface; Riemannian manifold ; The first eigenvalue ; Unit sphere

1 Introduction and main results

Let M^n be an embedded compact orientable minimal hypersurface in an $(n+1)$-dimensional compact orientable Riemannian manifold N.

In 1982, Yau [4] proposed the following conjecture.

Yau' Conjecture. *If $\lambda_1(M^n)$ is the first eigenvalue of an embedded compact minimal hypersurface M^n of unit sphere $\mathbb{S}^{n+1}(1)$, the standard $(n+1)$-sphere of sectional curvature 1, then $\lambda_1(M^n)=n$.*

In 1983, Choi-Wang [1] showed that if M^n is an embedded compact minimal hypersurface of unit sphere $\mathbb{S}^{n+1}(1)$, then $\lambda_1(M^n)\geqslant\frac{n}{2}$, where $\lambda_1(M^n)$ is the first eigenvalue of the Laplacian of M^n.

In the paper, in order to prove Yau' Conjecture, first we obtain the following result in Riemannian manifold N.

Theorem 1.1. *Let M^n be an embedded compact orientable minimal hypersurface in an $(n+1)$-dimensional compact orientable Riemannian manifold N. Suppose that $\lambda_1(M^n)$ is the first eigenvalue of the Laplacian of M^n. If the Ricci curvature of N is bounded below by a positive constant k, then*

$$\frac{k}{2}+\frac{n}{2}\delta_1(u)\leqslant\lambda_1(M^n)\leqslant\frac{k}{2}+\frac{n}{2}\delta_2(u),$$

where

$$0<\delta_1(u)=\frac{\int_{\Omega_1}|\nabla u|^2\left(1-\sqrt{1-\frac{(n+1)k}{n}\frac{\int_{\Omega_1}u^2}{\int_{\Omega_1}|\nabla u|^2}}\right)^2}{(n+1)\int_{\Omega_1}u^2}\leqslant\frac{k}{n},$$

$$\delta_2(u)=\frac{\int_{\Omega_1}|\nabla u|^2\left(1+\sqrt{1-\frac{(n+1)k}{n}\frac{\int_{\Omega_1}u^2}{\int_{\Omega_1}|\nabla u|^2}}\right)^2}{(n+1)\int_{\Omega_1}u^2}\geqslant\frac{k}{n},$$

2010 *Mathematics Subject Classification.* Primary 35P15; Secondary 53C40, 53C42.
This work was supported by the National Natural Science Foundation of China (No. 11501500).
School of Mathematics Science, Huaibei Normal University, Huaibei, Anhui, 235000, China
E-mail: sunzhongyang12@163.com

and u is the solution of the Dirichlet problem such that

$$\begin{cases} \Delta u = 0, & \text{in } \Omega_1; \\ u = f, & \text{in } \partial\Omega_1 = M^n, \end{cases}$$

which is called Dirichlet problem (1).

We see that $0 < \delta_1(u) \leqslant \frac{k}{n}$, $\delta_2(u) \geqslant \frac{k}{n}$ in Theorem 1.1. In particular, we have $\delta_1(u) = \delta_2(u) = \frac{k}{n}$ iff $\int_{\Omega_1} |\nabla u|^2 = \frac{(n+1)k}{n} \int_{\Omega_1} u^2$. When we take $\int_{\Omega_1} |\nabla u|^2 = \frac{(n+1)k}{n} \int_{\Omega_1} u^2$ in Theorem 1.1, we have $\lambda_1(M^n) = k$. Note that the Ricci curvature of unit sphere $\mathbb{S}^{n+1}(1)$ is n and an embedded compact hypersurface of unit sphere $\mathbb{S}^{n+1}(1)$ is already orientable. We obtain Corollary 1.2 which partially solves Yau' Conjecture.

Corollary 1.2. *Let M^n be an embedded compact minimal hypersurface of unit sphere $\mathbb{S}^{n+1}(1)$. Suppose that $\lambda_1(M^n)$ is the first eigenvalue of the Laplacian of M^n. If the solution u of* Dirichlet problem (1) *satisfies $\int_{\Omega_1} |\nabla u|^2 = (n+1) \int_{\Omega_1} u^2$, then*

$$\lambda_1(M^n) = n.$$

According to (3.3) in the proof of Theorem 1.1, we know that $\int_{\Omega_1} |\nabla u|^2 \geqslant \frac{(n+1)k}{n} \int_{\Omega_1} u^2$ is naturally true and meaningful in Theorem 1.1. Note that the Ricci curvature of unit sphere $\mathbb{S}^{n+1}(1)$ is n. Hence, we get the following corollary.

Corollary 1.3. *Let M^n be an embedded compact minimal hypersurface of unit sphere $\mathbb{S}^{n+1}(1)$. Suppose that $\lambda_1(M^n)$ is the first eigenvalue of the Laplacian of M^n. Then*

$$\int_{\Omega_1} |\nabla u|^2 \geqslant (n+1) \int_{\Omega_1} u^2,$$

where u is the solution of Dirichlet problem (1).

Remark 1.4. Corollary 1.3 proves that the condition $\int_{\Omega_1} |\nabla u|^2 = (n+1) \int_{\Omega_1} u^2$ is naturally true and meaningful in Corollary 1.2.

From Lemma 2.1 and $\int_{\partial\Omega_1} h(\overline{\nabla} u, \overline{\nabla} u) \geqslant 0$ in the proof of Theorem 1.1, we can conclude the following corollary.

Corollary 1.5. *Let M^n be an embedded compact orientable minimal hypersurface in an $(n+1)$-dimensional compact orientable Riemannian manifold N. Suppose that $\lambda_1(M^n)$ is the first eigenvalue of the Laplacian of M^n. If the Ricci curvature of N is bounded below by a positive constant k, then*

$$\lambda_1(M^n) > \frac{k}{2}.$$

Note that the Ricci curvature of unit sphere $\mathbb{S}^{n+1}(1)$ is n. Hence, from Corollary 1.5 we obtain Corollary 1.6 which improves the result $\lambda_1(M^n) \geqslant \frac{n}{2}$ of [1].

Corollary 1.6. *Let M^n be an embedded compact minimal hypersurface of unit sphere $\mathbb{S}^{n+1}(1)$. Suppose that $\lambda_1(M^n)$ is the first eigenvalue of the Laplacian of M^n. Then*

$$\lambda_1(M^n) > \frac{n}{2}.$$

Finally, by Corollaries 1.2 and 1.3, what we want to know is whether the following problem is true.

Problem 1.7. *Let M^n be an embedded compact minimal hypersurface of unit sphere $\mathbb{S}^{n+1}(1)$. Suppose that $\lambda_1(M^n)$ is the first eigenvalue of the Laplacian of M^n. If the solution u of* Dirichlet problem (1) *satisfies $\int_{\Omega_1}|\nabla u|^2 > (n+1)\int_{\Omega_1} u^2$, then $\lambda_1(M^n) = n$?*

Remark 1.8. If Problem 1.7 is true, combining Corollary 1.2, Corollary 1.3 and Problem 1.7, then Yau' Conjecture is true.

2 Preliminaries

Let Ω_1 is a Riemannian manifold of dimensional $(n+1)$ with smooth boundary $\partial\Omega_1 = M^n$. Let u be a function defined on Ω_1 which is smooth up to $\partial\Omega_1$. The symbols Δu and ∇u will be respectively the Laplacian and the gradient of u with respect to the induced Riemannian metric on Ω_1 while $\overline{\Delta} u$ and $\overline{\nabla} u$ will be the Laplacian the gradient of u (defined on $\partial\Omega_1$) with respect to the induced Riemannian metric on $\partial\Omega_1$. For $x \in \Omega_1$ and $X, Y \in T_x\Omega_1$, we can define the Hessian tensor $(D^2u)(X,Y) = X(Yu) - (\nabla_X Y)u$, where $\nabla_X Y$ is the coviariant derivative of the Riemannian connection of Ω_1. The covariant derivative of the Riemannian connection of $\partial\Omega_1 = M^n$ is given by $\overline{\nabla}_X Y$.

Suppose that $\{e_1, e_2, \cdots, e_n, e_{n+1}\}$ is a local orthonormal frame such that at $x \in \partial\Omega_1$, $e_1, e_2, \cdots, e_n$, are tangent to $\partial\Omega_1$ and e_{n+1} is the outward normal vector. Let h be the second fundamental form, $h(\upsilon, \omega) = < \nabla_\upsilon e_n, \omega >$, where υ, ω are vectors tangent to $\partial\Omega_1 = M^n$, and H be the mean curvature, i.e., $H = \sum_{i=1}^n \frac{h(e_i,e_i)}{n}$.

We need the following lemmas will play a crucial role in the proof of Theorem 1.1.

Lemma 2.1. *Let M^n be an embedded compact orientable minimal hypersurface in an $(n+1)$-dimensional compact orientable Riemannian manifold N. Suppose that $\lambda_1(M^n)$ is the first eigenvalue of the Laplacian of M^n. If the Ricci curvature of N is bounded below by a positive constant k, then*

$$\big(2\lambda_1(M^n) - k\big)\int_{\Omega_1}|\nabla u|^2 - \int_{\partial\Omega_1} h(\overline{\nabla} u, \overline{\nabla} u) > 0,$$

where u is the solution of Dirichlet problem (1).

Proof. Since the Ricci curvature of N is strictly positive, the first Betti number of N must be zero. Since M^n and N are orientable, by looking at the exact sequences of homology groups, we can see that M^n divides N into components Ω_1 and Ω_2 such that $\partial\Omega_1 = \partial\Omega_2 = M^n$.

Let f be the first eigenfunction of M^n, i.e.,

$$\overline{\Delta} f + \lambda_1(M^n) f = 0. \tag{2.1}$$

Let u be the solution of the Dirichlet problem such that

$$\begin{cases} \Delta u = 0, & \text{in } \Omega_1; \\ u = f, & \text{in } \partial\Omega_1 = M^n, \end{cases} \tag{2.2}$$

which is called Dirichlet problem (1).

So u is a function defined on Ω_1 smooth up to $\partial\Omega_1$. Then we have

$$\Delta u = \sum_{i=1}^{n+1} D^2u(e_i, e_i) = \sum_{i=1}^{n+1} u_{ii}, \tag{2.3}$$

where $u_{ij} = D^2u(e_i, e_j), \qquad i, j = 1, \cdots, n+1.$

When $i \neq n+1$ and $x \in \partial\Omega_1$, we get

$$\nabla_{e_i} e_i = \overline{\nabla}_{e_i} e_i - h_{ii} e_{n+1}, \tag{2.4}$$

where $h_{ij} = h(e_i, e_j)$.

Hence when $x \in \partial\Omega_1 = M^n$, we can conclude from (2.2), (2.3) and (2.4) that

$$\Delta u = u_{(n+1)(n+1)} + \overline{\Delta} f + \sum_{i=1}^{n} h_{ii} e_{n+1}(u) = u_{(n+1)(n+1)} + \overline{\Delta} f + nHu_{n+1}, \tag{2.5}$$

where $u_{n+1} =< \nabla u, e_{n+1} >$ and H is the mean curvature of M^n.

For $x \in \partial\Omega_1$, we know from (2.1) and (2.5) that

$$u_{(n+1)(n+1)} = \lambda_1(M^n) f - nHu_{n+1}. \tag{2.6}$$

For $x \in \Omega_1$, note that the fact $\Delta|\nabla u|^2 = 2\sum_{i,j=1}^{n+1} u_{ij}^2 + 2\sum_{i,j=1}^{n+1} R_{ij}u_iu_j + 2\sum_{i=1}^{n+1} u_i(\Delta u)_i$ can be found in [2]. Since $\Delta u = 0$, we have

$$\Delta|\nabla u|^2 = 2\sum_{i,j=1}^{n+1} u_{ij}^2 + 2\sum_{i,j=1}^{n+1} R_{ij}u_iu_j, \tag{2.7}$$

where $R_{ij} = \mathrm{Ric}(e_i, e_j)$.

Since Ricci curvature of N is bounded by k, by (2.7) we have

$$\Delta|\nabla u|^2 \geqslant 2|D^2u|^2 + 2k|\nabla u|^2. \tag{2.8}$$

Then integrating (2.8) we have

$$\int_{\Omega_1} \Delta|\nabla u|^2 \geqslant 2\int_{\Omega_1} |D^2u|^2 + 2k\int_{\Omega_1} |\nabla u|^2. \tag{2.9}$$

When $i \neq n+1$, we have

$$u_{i(n+1)} = D^2u(e_i, e_{n+1}) = e_i(e_{n+1}u) - (\nabla_{e_i} e_{n+1})u = e_i(u_{n+1}) - \sum_{j=1}^{n} h_{ij}u_j. \tag{2.10}$$

By using the Stokes theorem, we can conclude from (2.1),(2.6) and (2.10) that

$$\begin{aligned}
\int_{\Omega_1} \Delta|\nabla u|^2 =& 2\int_{\partial\Omega_1} \sum_{i=1}^{n} u_i u_{i(n+1)} + 2\int_{\partial\Omega_1} u_{n+1}u_{(n+1)(n+1)} \\
=& 2\int_{\partial\Omega_1} \overline{\nabla} f \cdot \overline{\nabla} u_{n+1} - 2\int_{\partial\Omega_1} \sum_{i,j=1}^{n} h_{ij}u_iu_j + 2\int_{\partial\Omega_1} u_{n+1}u_{(n+1)(n+1)} \\
=& -2\int_{\partial\Omega_1} u_{n+1}\overline{\Delta} f - 2\int_{\partial\Omega_1} h(\overline{\nabla} u, \overline{\nabla} u) + 2\int_{\partial\Omega_1} u_{n+1}u_{(n+1)(n+1)} \\
=& 4\lambda_1(M^n)\int_{\partial\Omega_1} u_{n+1} f - 2\int_{\partial\Omega_1} h(\overline{\nabla} u, \overline{\nabla} u) - 2n\int_{\partial\Omega_1} Hu_{n+1}^2.
\end{aligned} \tag{2.11}$$

Combining (2.2) and Stokes theorem, we have

$$\int_{\Omega_1} |\nabla u|^2 = -\int_{\Omega_1} u\Delta u + \int_{\partial\Omega_1} uu_{n+1} = \int_{\partial\Omega_1} u_{n+1} f. \tag{2.12}$$

From (2.11) and (2.12), we have

$$\int_{\Omega_1} \Delta|\nabla u|^2 = 4\lambda_1(M^n)\int_{\Omega_1} |\nabla u|^2 - 2\int_{\partial\Omega_1} h(\overline{\nabla} u, \overline{\nabla} u) - 2n\int_{\partial\Omega_1} Hu_{n+1}^2. \tag{2.13}$$

Since M^n is minimal, by (2.9) and (2.13), we have

$$\big(2\lambda_1(M^n) - k\big)\int_{\Omega_1} |\nabla u|^2 \geqslant \int_{\partial\Omega_1} h(\overline{\nabla} u, \overline{\nabla} u) + \int_{\Omega_1} |D^2 u|^2. \tag{2.14}$$

We claim that

$$\int_{\Omega_1} |D^2 u|^2 \neq 0. \tag{2.15}$$

Otherwise, for all $1 \leqslant i, j \leqslant n+1$, we have $u_{ij} = 0$ on Ω_1. Since u is smooth up to $\partial\Omega_1$, for all $1 \leqslant i, j \leqslant n$, we have $f_{ij} = 0$ on M^n, which implies that $\overline{\Delta} f = 0$ which is impossible since f is the first eigenfunction of M^n. Thus, our claim is true. Hence from (2.15) we get

$$\int_{\Omega_1} |D^2 u|^2 > 0. \tag{2.16}$$

From (2.14) and (2.16), we have

$$\big(2\lambda_1(M^n) - k\big)\int_{\Omega_1} |\nabla u|^2 - \int_{\partial\Omega_1} h(\overline{\nabla} f, \overline{\nabla} f) > 0. \tag{2.17}$$

This completes the proof of Lemma 2.1. □

Lemma 2.2. *Let M^n be an embedded compact orientable minimal hypersurface in an $(n+1)$-dimensional compact orientable Riemannian manifold N. Suppose that $\lambda_1(M^n)$ is the first eigenvalue of the Laplacian of M^n. If the Ricci curvature of N is bounded below by a positive constant k, then*

$$\begin{aligned} Q(t) = &\left[\big(2\lambda_1(M^n) - k\big)\int_{\Omega_1} |\nabla u|^2 - \int_{\partial\Omega_1} h(\overline{\nabla} f, \overline{\nabla} f)\right]\cdot t^2 + 2\lambda_1(M^n)\int_{\Omega_1} u^2 \cdot t \\ &+ \frac{n}{n+1}\int_{\Omega_1} u^2 \geqslant 0, \qquad \forall\ t \in \mathbb{R}, \end{aligned} \tag{2.18}$$

where u is the solution of Dirichlet problem (1).

Proof. Let $z = \frac{\partial u}{\partial \nu} = u_\nu$ be the normal outward derivative of u. We know that the Reilly formula:

$$\int_{\Omega_1} (\Delta u)^2 = \int_{\Omega_1} |D^2 u|^2 + \int_{\Omega_1} \mathrm{Ric}(\nabla u, \nabla u) + \int_{\partial\Omega_1} 2z\overline{\Delta} u + \int_{\partial\Omega_1} h(\overline{\nabla} u, \overline{\nabla} u) + \int_{\partial\Omega_1} nHz^2. \tag{2.19}$$

Case 1: $t = 0$, then (2.18) is naturally true. Case 2: $t \neq 0$, we will consider the following Dirichlet problem

$$\begin{cases} \Delta g = u, & \text{in } \Omega_1; \\ g = tf, & \text{in } \partial\Omega_1 = M^n. \end{cases} \tag{2.20}$$

Combining Green formula and (2.20), we have

$$\begin{cases} \int_{\partial\Omega_1} f\frac{\partial u}{\partial \nu} = \int_{\Omega_1} |\nabla u|^2; \\ t\int_{\partial\Omega_1} f\frac{\partial u}{\partial \nu} = \int_{\Omega_1} < \nabla u, \nabla g >; \\ \int_{\partial\Omega_1} f\frac{\partial g}{\partial \nu} = \int_{\Omega_1} u^2 + \int_{\Omega_1} < \nabla u, \nabla g > . \end{cases} \tag{2.21}$$

From (2.21), we have

$$\int_{\Omega_1} < \nabla u, \nabla g >= t \int_{\Omega_1} |\nabla u|^2. \tag{2.22}$$

From (2.22) and Cauchy-Schwarz inequality, we have

$$\int_{\Omega_1} |\nabla g|^2 \geqslant t^2 \int_{\Omega_1} |\nabla u|^2. \tag{2.23}$$

From the third equation in (2.21) and (2.22), we get

$$t \int_{\partial\Omega_1} f \frac{\partial g}{\partial \nu} = t \int_{\Omega_1} u^2 + t^2 \int_{\Omega_1} |\nabla u|^2. \tag{2.24}$$

Since M^n is minimal and $|D^2 g|^2 \geqslant \frac{1}{n+1}(\Delta g)^2$, applying (2.19) to g, we obtain

$$\frac{n}{n+1} \int_{\Omega_1} (\Delta g)^2 \geqslant k \int_{\Omega_1} |\nabla g|^2 + 2 \int_{\partial\Omega_1} \frac{\partial g}{\partial \nu} \overline{\Delta}(tf) + \int_{\partial\Omega_1} h(\overline{\nabla} g, \overline{\nabla} g). \tag{2.25}$$

On the other hand, combining (2.1), (2.23), (2.24), (2.25) and $\Delta g = u$, we have

$$\frac{n}{n+1} \int_{\Omega_1} u^2 \geqslant kt^2 \int_{\Omega_1} |\nabla u|^2 - 2\lambda_1(M^n) \left[t \int_{\Omega_1} u^2 + t^2 \int_{\Omega_1} |\nabla u|^2 \right] + t^2 \int_{\partial\Omega_1} h(\overline{\nabla} f, \overline{\nabla} f). \tag{2.26}$$

Hence, we have

$$\begin{aligned} Q(t) = &\left[\big(2\lambda_1(M^n) - k\big) \int_{\Omega_1} |\nabla u|^2 - \int_{\partial\Omega_1} h(\overline{\nabla} f, \overline{\nabla} f) \right] \cdot t^2 + 2\lambda_1(M^n) \int_{\Omega_1} u^2 \cdot t \\ &+ \frac{n}{n+1} \int_{\Omega_1} u^2 \geqslant 0, \qquad \forall\ t \in \mathbb{R}. \end{aligned}$$

This completes the proof of Lemma 2.2. □

3 Proof of Theorem 1.1

Proof of Theorem 1.1. Combining (2.17) and (2.18), we know that the discriminant of $Q(t)$ is non-positive, i.e.,

$$\frac{n+1}{n} \lambda_1^2(M^n) \frac{\int_{\Omega_1} u^2}{\int_{\Omega_1} |\nabla u|^2} - 2\lambda_1(M^n) + k + \frac{\int_{\partial\Omega_1} h(\overline{\nabla} f, \overline{\nabla} f)}{\int_{\Omega_1} |\nabla u|^2} \leqslant 0. \tag{3.1}$$

Since $\int_{\partial\Omega_1} h(\overline{\nabla} u, \overline{\nabla} u) = \int_{M^n} h(\overline{\nabla} f, \overline{\nabla} f)$ and the outward normal vector of $\partial\Omega_2$ is $-e_n$, we have

$$\int_{\partial\Omega_2} h(\overline{\nabla} u, \overline{\nabla} u) = - \int_{\partial\Omega_1} h(\overline{\nabla} u, \overline{\nabla} u).$$

Hence we can assume that $\int_{\partial\Omega_1} h(\overline{\nabla} u, \overline{\nabla} u) \geqslant 0$; otherwise, we work with Ω_2 rather with Ω_1. Since $\int_{\partial\Omega_1} h(\overline{\nabla} f, \overline{\nabla} f) \geqslant 0$, from (3.1) we have

$$\frac{n+1}{n} \lambda_1^2(M^n) \frac{\int_{\Omega_1} u^2}{\int_{\Omega_1} |\nabla u|^2} - 2\lambda_1(M^n) + k \leqslant 0. \tag{3.2}$$

Combining (2.17), $\int_{\partial\Omega_1} h(\overline{\nabla} f, \overline{\nabla} f) \geqslant 0$ and $\int_{\Omega_1} |\nabla u|^2 > 0$, we have $\lambda_1(M^n) > \frac{k}{2}$. When we take $t = -\frac{n}{(n+1)k}$ in (2.18), combining $\lambda_1(M^n) > \frac{k}{2}$, $\int_{\partial\Omega_1} h(\overline{\nabla} f, \overline{\nabla} f) \geqslant 0$ and $\int_{\Omega_1} |\nabla u|^2 > 0$, we get

$$1 - \frac{(n+1)k}{n} \frac{\int_{\Omega_1} u^2}{\int_{\Omega_1} |\nabla u|^2} \geqslant 0. \tag{3.3}$$

From (3.2) and (3.3), we have

$$\begin{aligned}\lambda_1(M^n) \geqslant & \frac{n\int_{\Omega_1}|\nabla u|^2 - n\int_{\Omega_1}|\nabla u|^2\sqrt{1-\frac{(n+1)k}{n}\frac{\int_{\Omega_1}u^2}{\int_{\Omega_1}|\nabla u|^2}}}{(n+1)\int_{\Omega_1}u^2} \\ = & \frac{k(n+1)\int_{\Omega_1}u^2}{2n(n+1)\int_{\Omega_1}u^2} \\ & + \frac{2n\int_{\Omega_1}|\nabla u|^2 - k(n+1)\int_{\Omega_1}u^2 - 2n\int_{\Omega_1}|\nabla u|^2\sqrt{1-\frac{(n+1)k}{n}\frac{\int_{\Omega_1}u^2}{\int_{\Omega_1}|\nabla u|^2}}}{2(n+1)\int_{\Omega_1}u^2} \\ = & \frac{k}{2} + \frac{n}{2}\delta_1(u)\end{aligned} \tag{3.4}$$

and

$$\begin{aligned}\lambda_1(M^n) \leqslant & \frac{n\int_{\Omega_1}|\nabla u|^2 + n\int_{\Omega_1}|\nabla u|^2\sqrt{1-\frac{(n+1)k}{n}\frac{\int_{\Omega_1}u^2}{\int_{\Omega_1}|\nabla u|^2}}}{(n+1)\int_{\Omega_1}u^2} \\ = & \frac{k(n+1)\int_{\Omega_1}u^2}{2n(n+1)\int_{\Omega_1}u^2} \\ & + \frac{2n\int_{\Omega_1}|\nabla u|^2 - k(n+1)\int_{\Omega_1}u^2 + 2n\int_{\Omega_1}|\nabla u|^2\sqrt{1-\frac{(n+1)k}{n}\frac{\int_{\Omega_1}u^2}{\int_{\Omega_1}|\nabla u|^2}}}{2(n+1)\int_{\Omega_1}u^2} \\ = & \frac{k}{2} + \frac{n}{2}\delta_2(u),\end{aligned} \tag{3.5}$$

where

$$\delta_1(u) = \frac{\int_{\Omega_1}|\nabla u|^2\left(1-\sqrt{1-\frac{(n+1)k}{n}\frac{\int_{\Omega_1}u^2}{\int_{\Omega_1}|\nabla u|^2}}\right)^2}{(n+1)\int_{\Omega_1}u^2} \tag{3.6}$$

and

$$\delta_2(u) = \frac{\int_{\Omega_1}|\nabla u|^2\left(1+\sqrt{1-\frac{(n+1)k}{n}\frac{\int_{\Omega_1}u^2}{\int_{\Omega_1}|\nabla u|^2}}\right)^2}{(n+1)\int_{\Omega_1}u^2}. \tag{3.7}$$

Finally we will prove $0 < \delta_1(u) \leqslant \frac{k}{n}$, $\delta_2(u) \geqslant \frac{k}{n}$. Firstly, from (3.6) we have $\delta_1(u) \geqslant 0$. But $\delta_1(u) \neq 0$, otherwise, from (3.6) we can conclude that $1-\sqrt{1-\frac{(n+1)k}{n}\frac{\int_{\Omega_1}u^2}{\int_{\Omega_1}|\nabla u|^2}} = 0$, which implies that $\int_{\Omega_1}u^2 = 0$ which is impossible. Hence we get $\delta_1(u) > 0$ and $1-\frac{(n+1)k}{n}\frac{\int_{\Omega_1}u^2}{\int_{\Omega_1}|\nabla u|^2} \neq 1$. Combining (3.3), we know that

$$0 \leqslant 1 - \frac{(n+1)k}{n}\frac{\int_{\Omega_1}u^2}{\int_{\Omega_1}|\nabla u|^2} < 1. \tag{3.8}$$

Obviously, $\delta_1(u)$ can be written as

$$\delta_1(u) = \frac{2\int_{\Omega_1}|\nabla u|^2\left(1-\frac{(n+1)k}{n}\frac{\int_{\Omega_1}u^2}{\int_{\Omega_1}|\nabla u|^2} - \sqrt{1-\frac{(n+1)k}{n}\frac{\int_{\Omega_1}u^2}{\int_{\Omega_1}|\nabla u|^2}}\right) + \frac{(n+1)k}{n}\int_{\Omega_1}u^2}{(n+1)\int_{\Omega_1}u^2}. \tag{3.9}$$

From (3.8), we obtain

$$1 - \frac{(n+1)k}{n}\frac{\int_{\Omega_1}u^2}{\int_{\Omega_1}|\nabla u|^2} - \sqrt{1-\frac{(n+1)k}{n}\frac{\int_{\Omega_1}u^2}{\int_{\Omega_1}|\nabla u|^2}} \leqslant 0. \tag{3.10}$$

We can conclude from (3.9) and (3.10) that

$$\delta_1(u) \leqslant \frac{k}{n}.$$

Secondly, from (3.7) and (3.8) we obtain

$$\begin{aligned}\delta_2(u) =& \frac{\int_{\Omega_1} |\nabla u|^2 \left(1 + \sqrt{1 - \frac{(n+1)k}{n} \frac{\int_{\Omega_1} u^2}{\int_{\Omega_1} |\nabla u|^2}}\right)^2}{(n+1)\int_{\Omega_1} u^2} \\ =& \frac{2\int_{\Omega_1} |\nabla u|^2 \left(1 - \frac{(n+1)k}{n} \frac{\int_{\Omega_1} u^2}{\int_{\Omega_1} |\nabla u|^2} + \sqrt{1 - \frac{(n+1)k}{n} \frac{\int_{\Omega_1} u^2}{\int_{\Omega_1} |\nabla u|^2}}\right) + \frac{(n+1)k}{n}\int_{\Omega_1} u^2}{(n+1)\int_{\Omega_1} u^2} \\ \geqslant& \frac{k}{n}.\end{aligned}$$

So the proof of Theorem 1.1 is finished.

□

4 Applications

In 1980, Yang and Yau [3] obtained the following result.

Theorem 4.1 ([3]). *If M^2 is an orientable Riemannian surface of genus* g *with area A, then*

$$\lambda_1(M^2) \leqslant 8\pi(\mathrm{g}+1)A^{-1}.$$

From Corollary 1.5 and Theorem 4.1, we obtain the following theorem.

Theorem 4.2. *Let M^2 be an embedded compact orientable minimal surface of genus* g *with area A in a 3-dimensional compact orientable Riemannian manifold N. If the Ricci curvature of N is bounded below by a positive constant k, then*

$$A < \frac{16\pi(\mathrm{g}+1)}{k}.$$

Note that the Ricci curvature of unit sphere $\mathbb{S}^{n+1}(1)$ is n. From Theorem 4.2, we get the following corollary.

Corollary 4.3. *Let M^2 be an embedded compact minimal surface of genus* g *with area A in unit sphere $\mathbb{S}^3(1)$. Then*

$$A < 8\pi(\mathrm{g}+1).$$

From Corollary 1.2 and Theorem 4.1, we get the following theorem.

Theorem 4.4. *Let M^2 be an embedded compact minimal surface of genus* g *with area A in unit sphere $\mathbb{S}^3(1)$. If the solution u of* Dirichlet problem (1) *satisfies $\int_{\Omega_1} |\nabla u|^2 = 3\int_{\Omega_1} u^2$, then*

$$A \leqslant 4\pi(\mathrm{g}+1).$$

Naturally, we propose the following conjecture.

Conjecture 4.5. *Let M^2 be an embedded compact minimal surface of genus* g *with area A in unit sphere $\mathbb{S}^3(1)$. Then*

$$A \leqslant 4\pi(\mathrm{g}+1).$$

References

[1] H. I. Choi and A. N. Wang, A first eigenvalue estimate for minimal hypersurfaces, J. Differential Geom., **18**(1983) 559-562.

[2] R. Schoen and S.T. Yau, Lectures on Differential Geometry. Cambridge, MA, 1994.

[3] P. Yang and S. T. Yau, Eigenvalue of the Laplacian of compact Riemann surfaces and minimal submanifolds, Ann. Scuola Norm. Sup. Pisa, **7**(1980) 55-63.

[4] S. T. Yau, Problem section, Seminar on differential geometry, Annals of Math. Studies, No. 102, Princeton Univ. Press, Princeton, N.J., 1982.